

## Five new species and a new combination in Cyperaceae from the Kimberley region of Western Australia

Barbara L. Rye<sup>1</sup>, Russell L. Barrett<sup>1,2,3,6,7</sup>, Matthew D. Barrett<sup>1,2,3</sup>, Jeremy J. Bruhl<sup>4</sup>,  
Kerri L. Clarke<sup>4</sup> and Karen L. Wilson<sup>5</sup>

<sup>1</sup>Western Australian Herbarium, Department of Parks and Wildlife,  
Locked Bag 104, Bentley Delivery Centre, Western Australia 6983

<sup>2</sup>Botanic Gardens and Parks Authority, Kings Park and Botanic Garden, West Perth, Western Australia 6005

<sup>3</sup>School of Plant Biology, Faculty of Science, The University of Western Australia, Crawley, Western Australia 6009

<sup>4</sup>Botany, School of Environmental and Rural Science, University of New England, Armidale, New South Wales 2351

<sup>5</sup>National Herbarium of New South Wales, Royal Botanic Gardens and Domain Trust, Sydney, New South Wales 2000

<sup>6</sup>Current address: Australian National Herbarium, Centre for Australian National Biodiversity Research, National Research  
Collections Australia, CSIRO, GPO Box 1600, Canberra, Australian Capital Territory 2601

<sup>7</sup>Corresponding author, email: russell.barrett@csiro.au

### Abstract

Rye, B.L., Barrett, R.L., Barrett, M.D., Bruhl, J.J., Clarke, K.L. & Wilson, K.L. Five new species and a new combination in Cyperaceae from the Kimberley region of Western Australia. *Nuytsia* 26: 167–184 (2015). Four new species of *Actinoschoenus* Benth. and one new species of *Fimbristylis* Vahl are described from the north Kimberley region of Western Australia: *A. glabrispiculus* Rye, R.L.Barrett & M.D.Barrett, *A. pentagonus* Rye, R.L.Barrett & M.D.Barrett, *A. quadricostatus* Rye, R.L.Barrett & M.D.Barrett, *A. ramosus* Rye, R.L.Barrett & M.D.Barrett, and *F. helicophylla* Rye, R.L.Barrett & M.D.Barrett. One new combination is made: *A. arthrostyloides* (W.Fitzg.) K.L.Clarke, K.L.Wilson & J.J.Bruhl. Several of these species have very restricted distributions and are of conservation concern. Two species also occur in the western Northern Territory. All of these taxa have previously been listed on Western Australia's plant census under phrase names. All species are illustrated. A revised key to species of *Actinoschoenus* in the Kimberley region is provided.

### Introduction

Cyperaceae are well represented in the Kimberley region with about 17 genera and 190 species currently recorded. Several genera in the regional flora treatment by Rye (1992) contained informally named species, indicative of the extent of taxonomic uncertainties for the sedges of this region. Further collecting, field study and examination of type specimens has enabled us to establish formal names for the species of *Actinoschoenus* Benth. and one distinctive species of *Fimbristylis* Vahl.

Full descriptions for the five new species named here—*A. glabrispiculus* Rye, R.L.Barrett & M.D.Barrett, *A. pentagonus* Rye, R.L.Barrett & M.D.Barrett, *A. quadricostatus* Rye, R.L.Barrett & M.D.Barrett, *A. ramosus* Rye, R.L.Barrett & M.D.Barrett and *F. helicophylla* Rye, R.L.Barrett & M.D.Barrett—were published previously by Rye (1992) under informal names. A new combination is provided for *A. arthrostyloides* (W.Fitzg.) K.L.Clarke, K.L.Wilson & J.J.Bruhl based on *F. arthrostyloides* W.Fitzg. (Fitzgerald 1918). Four species are endemic to the Kimberley region of Western Australia, while

two extend into the adjacent Northern Territory. All of these species have previously been known by informal phrase names in Western Australia and four of them have conservation priority (Western Australian Herbarium 1998–).

### Methods

All measurements are based on dried herbarium material. Most species newly described here have been examined in the field by R.L. and M.D. Barrett. Dry leaves, culms and seeds were mounted on stubs using double-sided carbon tape with conductive carbon paint, coated with gold using an EMITECH K550X Sputter Coater and imaged at high vacuum and high voltage (15 KV) using a Jeol JCM 6000 NeoScope bench-top Scanning Electron Microscope at Kings Park and Botanic Garden.

Photosynthetic pathway prediction was based on the ‘one cell distant criterion’ of Hattersley and Watson (1975) where 1) mesophyll chlorenchyma cells are not more than one chlorenchyma cell distant from the nearest vascular bundle, which predicts  $C_4$ , or 2) a count of more than one indicates  $C_3$  status (see Bruhl *et al.* 1987 for a more detailed explanation and application to Cyperaceae). Observations here were of culm anatomy (*cf.* Bruhl *et al.* 1987; Bruhl & Wilson 2008) for two to four collections of each species of *Actinoschoenus* treated in this paper (Table 1). Hand-cut transverse sections from the rehydrated mid-third region of culms were double-stained with 0.5% acidified Astra Blue and 0.125% Basic Fuchsin in 50% ethanol for *c.* 1–2 min and 2 s respectively. Sections were rinsed between stains and mounted in 50% glycerol for photomicroscopy. Images were captured with a Nikon DS-Ri1 digital camera on a motorised Nikon Eclipse 90i microscope using the ‘large image’ feature of NIS Elements AR3.22.00 software to capture and stitch multiple images.

**Table 1.** Vouchers used for photosynthetic pathway anatomical observations in *Actinoschoenus*. Asterisk (\*) specimens under *A. ramosus* are from the East Alligator River area, Northern Territory, and require further taxonomic study (see text).

<i>Actinoschoenus</i> species	Vouchers
<i>A. arthrotyloides</i>	<i>C.R. Dunlop</i> 5238 (PERTH 01345354)
	<i>K.F. Kenneally</i> 9948 (PERTH 01428888)
	<i>J. Russell-Smith</i> 7584 & <i>D. Lucas</i> (DNA 0037927)
<i>A. glabrispiculus</i>	<i>K.F. Kenneally</i> 4474 (CANB 292219)
	<i>K.F. Kenneally</i> 4789 (PERTH 01489054)
	<i>J.H. Willis s.n.</i> (PERTH 01489070)
<i>A. pentagonus</i>	<i>C.R. Dunlop</i> 5309 (CANB 293319)
	<i>K.F. Kenneally</i> 11266 (PERTH 02249154)
<i>A. quadricostatus</i>	<i>G.W. Holmes s.n.</i> (MEL 204869)
	<i>P.G. Wilson</i> 11426 (PERTH 01489011)
	<i>C.R. Dunlop</i> 4403 (DNA 0010763)*
<i>A. ramosus</i>	<i>C.R. Dunlop</i> 5303 (DNA 0017460)
	<i>R. Fensham</i> 872 (DNA 0054914)*
	<i>J. Russell-Smith</i> 8423 & <i>J. Brock</i> (DNA 0055610)*

## Anatomy and photosynthetic pathway

A detailed database of photosynthetic pathway in the family Cyperaceae has been developed covering almost all genera and around 25% of all species (Bruhl & Wilson 2008), including one out of about ten species of *Actinoschoenus*. All samples of *Actinoschoenus* examined in this study (Table 1) consistently exhibited a maximum cells-distant count of more than one, indicative of C<sub>3</sub> photosynthetic pathway (Figure 1). Bruhl *et al.* (1987) and Bruhl and Wilson (2008) assessed a variety of methods for determining whether Cyperaceae species had C<sub>3</sub> or C<sub>4</sub> photosynthetic pathways, including anatomical (using the ‘one cell distant criterion’), CO<sub>2</sub> compensation point analyses and δ<sup>13</sup>C values. Anatomical assessment accurately predicted photosynthetic pathways for all Cyperaceae, except in some species of *Eleocharis* R.Br. The results of the present anatomical study are consistent with such findings and, with 60% of species in the genus now having been studied, make it likely that all species of *Actinoschoenus* will have C<sub>3</sub> anatomy.

## Taxonomy

### *Actinoschoenus* Benth.

The circumscription of genera in Abildgaardieae Lye and ‘Arthrostyleidae’ is under investigation. *Actinoschoenus* is a member of the ‘Arthrostyleidae’ which is embedded within the tribe Abildgaardieae and sister to *Fimbristylis* (Hinchliff & Roalson 2013). *Actinoschoenus* was included in a more broadly circumscribed *Fimbristylis* by Latz (1990). Revision of the other Australian species distributed in the Northern Territory and possibly Queensland, will follow separately (Clarke *et al.*, unpubl. data). The treatment presented here is adapted from Rye (1992).

### Key to *Actinoschoenus* species in the Kimberley region of Western Australia based on Rye (1992)

1. Inflorescence of 3 or more spikelets, with 2–4 primary rays and often also secondary rays each bearing 1 spikelet. Uppermost 2 glumes 4.5–5 mm long ..... **A. ramosus**
- 1: Inflorescence a solitary spikelet or a head of 2–7 spikelets. Uppermost 2 glumes 6–9 mm long
  2. Inflorescence bracts 2–5, obvious, the basal bract longest. Culms densely hairy, with two types of hairs, most of the hairs short but some much longer ones scattered along the ridges
    3. Culms obtusely 5(6)-ridged. Spikelets (1–)2–7. Upper glumes distally minutely hairy on sides as well as midvein ..... **A. arthrostyleoides**
    - 3: Culms acutely 4-ridged. Spikelets 1–3. Upper glumes appearing glabrous at 10× or with a few hairs on midvein ..... **A. quadricostatus**
  - 2: Inflorescence bracts many, not obvious, intergrading with the glumes, the uppermost bract longest. Culms nearly glabrous to densely hairy, the hairs all short
    4. Culms 0.2–0.35 mm in diam., 4-ridged. Inflorescence of 1 spikelet; bracts and glumes glabrous except for 2 basal bracts. .... **A. glabrispiculus**
    - 4: Culms 0.5–1.0 mm in diam., 5-ridged. Inflorescence of 3–6 spikelets; bracts and glumes densely minutely hairy. .... **A. pentagonus**




Figure 1. Astra Blue-Basic Fuchsin stained culm cross-sections of *Actinoschoenus* species showing culm features and  $C_3$  photosynthetic pathway anatomy. A–C – *A. arthrostyloides*; D, E – *A. glabrispiculus*; F – *A. pentagonus*, showing (in F2) typical  $C_3$  anatomy with parenchymatous bundle sheath cells without chloroplasts (arrow), and three cell layers of palisade parenchyma (i.e. a maximum cells-distant count of 2; stars); G – *A. quadricostatus*; H – *A. ramosus*; I–K – *A. aff. ramosus*. Scale bar = 200  $\mu\text{m}$  (A–F1, G–K); 40  $\mu\text{m}$  (F2). Images from C.R. Dunlop 5238 (A), K.F. Kenneally 9948 (B), J. Russell-Smith 7584 & D. Lucas (C), K.F. Kenneally 4474 (D), J.H. Willis s.n. (E), K.F. Kenneally 11266 (F), P.G. Wilson 11426 (G), C.R. Dunlop 5303 (H), C.R. Dunlop 4403 (I), J. Russell-Smith 8423 & J. Brock (J) and R. Fensham 872 (K). Photomicrographs by K.L. Clarke.

**Actinoschoenus arthrotyloides** (W.Fitzg.) K.L. Clarke, K.L. Wilson & J.J. Bruhl, *comb. nov.*

*Basionym:* *Fimbristylis arthrotyloides* W.Fitzg., *J. & Proc. Roy. Soc. West. Austral.* 3: 121 (1918). *Lectotype* (designated here): Artesian Range, near Walcott Inlet, Western Australia, August 1905, *W.V. Fitzgerald* 1368 (*lecto*: PERTH 01031120; *isolecto*: BM 000990865 image seen, BRIAQ0340989 image seen, NSW 696918, NSW 696919, PERTH 01031104, PERTH 01031112).

*Actinoschoenus* sp. B, B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1037 (1992).

*Actinoschoenus* sp. B Kimberley Flora (G.J. Keighery 2649), G. Paczkowska & A.R. Chapman, *West. Austral. Fl.: Descr. Cat.*, p. 41 (2000); Western Australian Herbarium, in *FloraBase*, <http://florabase.dpaw.wa.gov.au/> [accessed 1 March 2014].

*Perennial*, caespitose sedge, 0.3–0.8 m high. *Culms* 0.5–1 mm diam., overall shape in TS ± terete but strongly and obtusely 5(6)-ridged, densely covered by short hairs and with much longer hairs scattered along the ridges. *Leaves* densely covered by short and long hairs; uppermost leaf 35–100 mm long, with a blade 6–11 mm long. *Inflorescence* a head of (1)2–7 somewhat spreading spikelets. *Bracts* 2–5, hairy, the longest bract basal; basal bract 7–10 mm long, with a blade 6–8 mm long. *Spikelets* strongly compressed, narrowly ovate in outline, 7–9 mm long, 1.5–3 mm wide. *Glumes* 5–7, distichous, with a green keel and pale brown sides; upper glumes narrowly ovate, 6–8 mm long, distally minutely hairy, acute, often apiculate. *Stamens* 3; anther 3.5–4 mm long. *Style* undivided for 4.3–5.5 mm; base narrowly triangular to triangular in outline, *c.* 1 mm long, 0.5 mm wide; stigmatic branches 3, about as long as undivided portion of style. *Nut* with a stipe or contracted base 0.6–1 mm long; body whitish to grey-brown, 2–2.5 mm long, 1.5–2 mm wide, transversely tuberculate-ridged. *C*<sub>3</sub> photosynthetic pathway inferred from anatomy. (Figures 1A–C, 2)

*Diagnostic characters.* *Culms* densely hairy, displaying 2 hair types, obtusely 5(6)-ridged. *Inflorescence* (1–)2–7 spikelets; bracts 2–5, the basal bract longest. *Upper glumes* 6–8 mm long, distally minutely hairy on sides as well as midvein.

*Other specimens examined.* WESTERN AUSTRALIA: 9 km NNW of Mt Agnes, 9 Jan. 2001, *M.D. Barrett* MDB 1175 (NSW, PERTH); large gully, 10.8 km NE of junction of Youwanjela Creek and Prince Regent River, Prince Regent Nature Reserve, 20 Jan. 2007, *R.L. Barrett & M.D. Barrett* RLB 3670 (CANB, PERTH); Youwanjela pavement 1, 26.7 km E of Kings Cascades; 12.4 km NNE of junction of Youwanjela Creek and Prince Regent River, Prince Regent Nature Reserve, 25 Jan. 2007, *R.L. Barrett & M.D. Barrett* RLB 3840 (CANB, NSW, PERTH); HR1 Site, Harding Range, 20 km WNW of Munja, N of Walcott Inlet, 13 Jan. 2010, *R.L. Barrett, M. Maier & P. Kendrick* RLB 6008 (NE, NSW, PERTH); Cypress Valley, on W side of Morgan River, 3 km E of Theda Station Homestead, 9 Mar. 2014, *R.L. Barrett* RLB 8866 (BM, BRI, CANB, DNA, K, MEL, NE, NSW, PERTH); Mitchell River, 22 Feb. 1980, *C.R. Dunlop* 5238 (DNA, NT, PERTH); near Gariyeli Creek, Prince Regent River Reserve, 24 Aug. 1974, *A.S. George* 12594 (CANB, PERTH); overlooking AAB pitline and along last section of trapline 2, Hidden Island, Buccaneer Archipelago, 18 June 1982, *A.J.M. Hopkins* BA 0243 (PERTH); track to Mitchell Falls, Mitchell Plateau, 31 May 1988, *S.W.L. Jacobs* 5804 & *P.G. Wilson* (NSW); Surveyors Pool, Mitchell Plateau, 17 Feb. 1980, *G.J. Keighery* 2649 (PERTH); Mt Daghish vine thicket, 19 June 1987, *G.J. Keighery & J.J. Alford s.n.* (NSW, PERTH); Hunter River, W Kimberley, 26 May 1987, *K.F. Kenneally* 9948 (PERTH); Boongaree Island, Prince Frederick Harbour, 5 July 1973, *P.G. Wilson* 11374 (PERTH). NORTHERN TERRITORY: headwaters of Lalngang Creek, 16 May 1994, *I.D. Cowie* 5066 & *N.G. Walsh* (CANB, DNA, MEL *n.v.*, NT); Spirit Hills Conservation Area,




Figure 2. *Actinoschoemus arthrostyloides*. A – habitat on sand over sandstone; B – habit; C – culms; D–F – inflorescences showing anthers and/or style branches; G – SEM of culm; H – SEM of glume; I – SEM of nut. Scale bars = 500  $\mu$ m (G); 1 mm (H, I). Images from R.L. Barrett RLB 8866 (A–F) and R.L. Barrett RLB 3840 (G–I). Photographs by R.L. Barrett.

Nancys Gorge, 17 Aug. 1996, *I.D. Cowie* 7114 (DNA, NSW); Keep River National Park, Spirit Hills area, c. 35 km SW of Bullo River Homestead, 23 Mar. 2009, *I.D. Cowie* 12307 (DNA, NSW); 44 km SE of Oenpelli, 13 June 1978, *P.K. Latz* 7764 (NT); 11 km S Legune Station, 5 Mar. 1989, *J. Russell-Smith* 7584 & *D. Lucas* (DNA); Bradshaw Military Training Area, c. 80 km NW Timber Creek, 2 Apr. 2007, *B.M. Stuckey* & *I.D. Cowie* 56 (DNA, PERTH).

**Phenology.** Flowers and fruits recorded in February and from May to August.

**Distribution and habitat.** Occurs in open woodland with shrubs on sand associated with sandstone, sometimes occurring on the edges of pools. From the Kimberley region, extending from Mitchell Plateau and Boongaree Island south to near Walcott Inlet, east to Bullo River in the Northern Territory.

*Conservation status.* Currently known from thirteen locations in Western Australia and six in the adjacent Northern Territory, and sufficiently widespread not to require listing as a priority species.

*Etymology.* The epithet is from the genus *Arthrostylis* R.Br. with the Greek termination *-oides* (like), in reference to the similar appearance of this species to that genus.

*Typification.* The lectotype is selected from seven duplicate sheets located by us. The primary sets of Fitzgerald's collections from names published by him in 1918 are usually duplicated at NSW and PERTH, so it is customary to choose a lectotype from one of these herbaria. The set at PERTH is stated to be the main material by Maiden (in Fitzgerald 1918), so this material is given preference. PERTH 01031120 is chosen as it contains mature material with nuts and has a label in Fitzgerald's script.

*Notes.* While this taxon has been known for some time to belong in *Actinoschoenus*, the relevant combination has not previously been made.

Most similar to *A. quadricostatus* and readily distinguished by the obtusely 5(6)-ridged culms (*vs* acutely 4-ridged) and (1)2–7 spikelets per head (*vs* 1–3).

The vernacular name of Hairy Actinoschoenus is suggested.

**Actinoschoenus glabrispiculus** Rye, R.L.Barrett & M.D.Barrett, *sp. nov.*

*Type:* west of Drysdale River, Drysdale River National Park, Western Australia [precise locality withheld for conservation reasons], 29 April 2008, *R.L. Barrett & M.D. Barrett* RLB 4895 (*holo:* PERTH 08046808; *iso:* CANB, DNA, K, NE, NSW).

*Actinoschoenus* sp. D, B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1039, Figure 310C (1992).

*Actinoschoenus* sp. D Kimberley Flora (K.F. Kenneally 4789), G. Paczkowska & A.R. Chapman, *West. Austral. Fl.: Descr. Cat.*, p. 41 (2000); Western Australian Herbarium, in *FloraBase*, <http://florabase.dpaw.wa.gov.au/> [accessed 1 March 2014].

*Illustration.* B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1038, Figure 310C (1992) [as *Actinoschoenus* sp. D].

*Perennial*, caespitose sedge, 0.3–0.6 m high. *Culms* 0.2–0.3 mm diam, obtusely 4-ridged, shortly hairy throughout or glabrescent especially along the ridges. *Leaves* densely shortly hairy to partially glabrous; uppermost leaf 30–40 mm long, with a blade up to 3 mm long. *Inflorescence* of 1 spikelet. *Bracts* 2–many but not clearly differentiated from the glumes, gradually increasing in size from base upwards; basal bracts 2–2.5 mm long, with a point or short blade 0.5–1 mm long, slightly hairy; upper bracts glabrous. *Spikelet* narrowly ovoid, 9–11 mm long, 1–2 mm diam. *Glumes* (and all but basal 2 inflorescence bracts) 10–12, 4-ranked or nearly so, with a green keel and whitish sides, becoming medium brown, glabrous; upper glumes narrowly ovate, 7–9 mm long, apiculate. *Stamens* 3; anther not seen. *Style* undivided for *c.* 7.3 mm; base narrowly triangular in outline, *c.* 1 mm long, 0.5 mm wide; stigmatic branches 3, apparently shorter than undivided part. *Nut* with a stipe or contracted base 1–1.6 mm long; body whitish to grey-brown, 2–2.5 mm long, 1.4–1.8 mm wide, transversely tuberculate-rugose or somewhat tuberculate-ridged. C<sub>3</sub> photosynthetic pathway inferred from anatomy. (Figures 1D, E, 3)




Figure 3. *Actinoschoenus glabrispiculus*. A–habit; B–compact plant base resprouting following fire; C–culms; D–inflorescence; E – SEM of culm; F – SEM of glume; G – SEM of nut. Scale bars = 500 µm (E); 1 mm (F, G). Images from R.L. Barrett & M.D. Barrett RLB 4895. Photographs by R.L. Barrett.

*Diagnostic characters.* Culms 0.2–0.3 mm diam., 4-ridged, nearly glabrous to densely hairy, the hairs all short. Inflorescence of 1 spikelet; bracts many or intergrading with the glumes, the uppermost bract longest; glabrous except for 2 basal bracts. Upper glumes 7–9 mm long, glabrous.

*Other specimens examined.* WESTERN AUSTRALIA: [localities withheld for conservation reasons] 22 May 1984, E.A. Chesterfield, S.J. Forbes & J.H. Willis EAC 319 (CANB, K n.v., L n.v., MO n.v., NSW, NT, PERTH); 19 May 2003, K. Coate 676 (PERTH); 22 Feb. 1980, C.R. Dunlop 5239 (DNA, NT, PERTH); 19 Aug. 1975, K.F. Kenneally 4474 (CANB, NT, PERTH); 13 June 1976, K.F. Kenneally 4789 (NT, PERTH 4 sheets); 28 May 1987, K.F. Kenneally 10017 (PERTH); 10 May 1986, P.K. Latz 10290 (DNA, NSW, NT); 11 Aug. 2008, C. Sgherza 8 (PERTH); 22 May 1984, J.H. Willis s.n. (MEL n.v., NSW, NT, PERTH).

*Phenology.* Flowers and fruits recorded in February and from May to August.

*Distribution and habitat.* Occurs on sandstone in open woodland and shrubland. Endemic to the Kimberley region, extending from Bigge Island and the Mitchell Plateau east to Napier Broome Bay and Drysdale River National Park.

*Conservation status.* *Actinoschoenus glabrispiculus* is listed by Jones (2014) as Priority Three under Department of Parks and Wildlife Conservation Codes for Western Australian Flora, under the name *A. sp. D Kimberley Flora* (K.F. Kenneally 4789). Known from nine collections from four broad locations in the north-west Kimberley. Probably more widespread but poorly collected.



*Etymology.* The epithet is from the Latin *glabri-* (glabrous-) and *spicula* (spikelet), in reference to the glabrous spikelets in this species.

*Notes.* Most similar to *A. pentagonus* and readily distinguished by the inflorescence having only a single spikelet (*vs* 3–6), 4-ridged culms (*vs* 5-ridged), and glabrous upper bracts and glumes (*vs* densely minutely hairy).

The vernacular name of Smooth Actinoschoenus is suggested.

**Actinoschoenus pentagonus** Rye, R.L.Barrett & M.D.Barrett, *sp. nov.*

*Type:* Carson Escarpment, 41 km south-east of new Theda Station Homestead, Drysdale River National Park, Western Australia, 29 April 2008, *R.L. Barrett & M.D. Barrett* RLB 4890 (*holo:* PERTH 08046727; *iso:* NE, NSW).

*Actinoschoenus* sp. E, B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1039, Figure 310D (1992).

*Actinoschoenus* sp. E Kimberley Flora (C.R. Dunlop 5309), G. Paczkowska & A.R. Chapman, *West. Austral. Fl.: Descr. Cat.*, p. 41 (2000); Western Australian Herbarium, in *FloraBase*, <http://florabase.dpaw.wa.gov.au/> [accessed 1 March 2014].

*Illustration.* B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1038, Figure 310D (1992) [as *Actinoschoenus* sp. E].

*Perennial*, caespitose sedge, 0.4–1 m high. *Culms* 0.5–0.8 mm diam., obtusely 5-ridged (rarely a few culms 4-ridged), densely shortly hairy throughout or glabrescent especially on the ridges. *Leaves* densely hairy; uppermost leaf 35–70 mm long, the blade 2–5 mm long. *Inflorescence* of 3–6 scarcely spreading spikelets, often resembling a solitary spikelet. *Bracts* many, greenish, gradually increasing in size from base upwards, densely shortly hairy; basal bract 3–5 mm long; uppermost bract *c.* 6 mm long, with a point or short blade up to 1.7 mm long. *Spikelets* compressed, narrowly ovate in outline, 7–11 mm long, *c.* 2 mm wide. *Glumes* 4–6, distichous to almost spirally arranged, pale to medium brown, densely minutely hairy; uppermost glumes narrowly ovate, 7–9 mm long, acute, often apiculate. *Stamens* 3; anther *c.* 4 mm long. *Style* undivided for 5.3–7 mm; base triangular or narrowly triangular in outline, 0.8–1.5 mm long, 0.5–0.8 mm wide; stigmatic branches 3, about as long as or somewhat longer than remainder of style. *Nut* with a stipe or basal contraction 0.8–1.5 mm long; body whitish or pale brown, *c.* 3 mm long, 1.6–2 mm wide, smooth to slightly tuberculate-rugose between the angles. C<sub>3</sub> photosynthetic pathway inferred from anatomy. (Figures 1F, 4)

*Diagnostic characters.* *Culms* 0.5–0.8 mm diam., 5-ridged, nearly glabrous to densely hairy, the hairs all short. *Inflorescence* of 3–6 spikelets; bracts many or intergrading with the glumes, the uppermost bract longest; minutely hairy. *Uppermost glumes* 7–9 mm long, densely minutely hairy.

*Other specimens examined.* WESTERN AUSTRALIA: near creek on Kalumburu road, 174.4 km by road N from junction with Gibb River to Ellenbrae road, 30 Apr. 1985, *T.E.H. Aplin et al.* 817 (PERTH); *c.* 10 km E of Purulba Massif, Prince Regent Nature Reserve, 2 Feb. 1999, *M.D. Barrett* 710 (PERTH); 6.7 km NE of Bachsten Creek campsite, Prince Regent Nature Reserve, 3 Feb. 1999, *M.D. Barrett* 747 (PERTH); Roe River [pavement 1] on mainland, 3 km ESE of Gertrude Cove, Kiska Island, 23 Apr. 2008, *R.L. Barrett & M.D. Barrett* RLB 4597 (PERTH); HR1 Site, Harding Range,




Figure 4. *Actinoschoenus pentagonus*. A – plant base with sand-binding roots; B – culms; C – inflorescence; D – SEM of culm; E – SEM of glume; F – SEM of nut. Scale bars = 500  $\mu\text{m}$  (D); 1 mm (E, F). Images from R.L. Barrett & M.D. Barrett RLB 4597 (A–C) and R.L. Barrett & M.D. Barrett RLB 3776 (D–E). Photographs by R.L. Barrett.

20 km WNW of Munja, N of Walcott Inlet, 13 Jan. 2010, R.L. Barrett, M. Maier & P. Kendrick RLB 6009 (BRI, CANB, MEL, NE, NSW, PERTH); Governor Island, Napier Broome Bay, 19 May 1986, E.A. Chesterfield, S.J. Forbes & J.H. Willis EAC 247 (MEL, NSW, NT n.v., PERTH); Mitchell Plateau, 26 Feb. 1980, C.R. Dunlop 5309 [distributed as 5306] (BRI n.v., CANB, DNA, K n.v., NSW, NT n.v., PERTH); c. 1.4 km S of Ila Point, West Governor Island, Napier Broome Bay, 19 May 1984, S.J. Forbes 2061 (MEL n.v., NT n.v., PERTH); base of Anjo Peninsula (between Napier Broome Bay and Vansittart Bay), 5 km N of West Bay landing, 22 May 1984, S.J. Forbes 2114 (CANB, L n.v., MEL, NSW, NT n.v., PERTH); Blyxa Creek, Prince Regent River Reserve, 19 Aug. 1974, A.S. George 12454 (NT n.v., PERTH 2 sheets); Mertens Creek, Mitchell Plateau, 31 May 1988, S.W.L. Jacobs & Peter G. Wilson SWLJ 5813 (DNA n.v., NSW, PERTH); adjacent to King George River above Falls, 7 June 1992, K.F. Kenneally 11266 (NSW, PERTH); Mitchell Falls walking track, Mitchell Plateau, 20 July 1988, M. Parris 9392 (CANB, PERTH); c. 1 km SW of Mitchell Plateau track on western side of King Edward River crossing, 2 July 2002, K.L. Wilson 10096 (EIU, GENT, NSW, NY, PERTH, PRE); 6 km N of Kalumburu on Pago road, 3 July 2002, K.L. Wilson 10103 (NE, NSW, PERTH); parking area at start of path to Mitchell Falls, 1 July 2002, K.L. Wilson 10092 (L, NSW, PERTH). NORTHERN TERRITORY: Keep River National Park, Spirit Hills section, 10 May 2008, I.D. Cowie 12099 (B n.v., DNA n.v., MO n.v., NSW); Spirit Hills section, Keep River National Park, 10 May 2008, I.D. Cowie 12104 (DNA n.v., NSW, PERTH); c. 19 km NNW of Bullo River homestead, Spirit Hills, Keep River National Park, 22 Mar. 2009, I.D. Cowie 12291 (DNA n.v., MO n.v., NSW, PERTH); Spirit Hills Conservation Reserve, c. 100 km N of Ranger Station, 13 May 2011, D.L. Lewis 1720 (CANB, DNA n.v., NSW).

*Phenology.* Flowers and fruits from February to May, also recorded in August.

*Distribution and habitat.* Grows in sand or gravelly soil on sandstone in open woodland with *Triodia* spp. From the north Kimberley region, between Mitchell Plateau, Prince Regent River Reserve, Governor Island and King Edward River, and just into the western Northern Territory near Keep River.

*Conservation status.* *Actinoschoenus pentagonus* appears to be widespread and is not currently threatened.

*Etymology.* The epithet is from the Greek *penta-* (five-) and *-gonus* (angled), in reference to the five-ridged culms in this species.

*Notes.* The base of the plant seems to be more resinous than in the other species of *Actinoschoenus* described here. Because the spikelets are closely associated and the inflorescence bracts are numerous, the head can easily be mistaken for a single spikelet.

*Actinoschoenus pentagonus* appears to be similar to the species referred to as *Fimbristylis* sp. G in Latz (1990) [= *A.* sp. Mount Brockman (R.C. Hinz 362); CHAH 2007–] but with fewer culm ribs (5 vs 5–9), larger spikelets (7–11 vs 4.5–7 mm long) and fewer of them per head (3–6 vs 3–10 per head). It is most similar to *A. glabrispiculus* and readily distinguished by the inflorescence having 3–6 spikelets (vs solitary), 5-ridged culms (vs 4-ridged), and densely minutely hairy upper bracts and glumes (vs glabrous).

There appear to be two variants of this species, the typical variant with densely hairy culms (Figure 4B), and a second variant with glabrous to sparsely hairy culms (e.g. *R.L. Barrett & M.D. Barrett* RLB3776; Figure 4D). Further investigation is required to determine whether there are other differences that may justify the recognition of two taxa.

The vernacular name of Five-angled *Actinoschoenus* is suggested.

***Actinoschoenus quadricostatus*** Rye, R.L.Barrett & M.D.Barrett, *sp. nov.*

*Type:* Uwins Island, Bonaparte Archipelago, Western Australia [precise locality withheld for conservation reasons], 6 June 2008, *M.N. Lyons* 6020 (*holo:* PERTH 08615497; *iso:* CANB, NE, NSW).

*Actinoschoenus* sp. C, B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1037, Figure 310B (1992).

*Actinoschoenus* sp. C Kimberley Flora (P.G. Wilson s.n. 12/5/72), G. Paczkowska & A.R. Chapman, *West. Austral. Fl.: Descr. Cat.*, p. 41 (2000); Western Australian Herbarium, in *FloraBase*, <http://florabase.dpaw.wa.gov.au/> [accessed 1 March 2014].

*Illustration.* B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1038, Figure 310B (1992) [as *Actinoschoenus* sp. C].

*Perennial*, caespitose sedge, 0.2–0.6 m high. *Culms* 0.5–0.75 mm diam., acutely 4-ridged, densely shortly hairy and with scattered much longer hairs to 0.6 mm long. *Leaves* densely covered by short and long hairs; uppermost leaf 25–65 mm long, with a blade 6–10 mm long. *Inflorescence* of 1 spikelet or a head of 2 or 3 somewhat spreading spikelets. *Bracts* 2 or 3, hairy, the longest bract basal; basal bract 6–10 mm long, with a blade 5–8 mm long. *Spikelets* strongly compressed, narrowly ovate in outline, 7–9 mm long, 1.5–3 mm wide. *Glumes* 4–6, distichous, with a green keel and brown sides;

upper glumes narrowly ovate, 6–8 mm long, glabrous or with a few hairs on midvein, apiculate. *Stamens* 3; anther not seen. *Style* undivided for 5–6 mm; base narrowly triangular in outline, 1.5–2 mm long, c. 0.5 mm wide; stigmatic branches 3, about as long as remainder of style. *Nut* with a stipe or contracted base 0.7–1 mm long; body 2.2–2.5 mm long, 1.8–2 mm wide, transversely tuberculate-ridged.  $C_3$  photosynthetic pathway inferred from anatomy. (Figures 1G, 5)

*Diagnostic characters.* *Culms* acutely 4-ridged, densely hairy, most of the hairs short but some much longer ones scattered along the ridges. *Inflorescence* of 1–3 spikelets; bracts 2 or 3, the basal bract longest. *Upper glumes* 6–8 mm long, glabrous or with a few hairs on midvein.

*Other specimens examined.* WESTERN AUSTRALIA: [localities withheld for conservation reasons] May 1943, *Rev. G.W. Holmes s.n.* (MEL, PERTH); 13 Feb. 2009, *G.J. Keighery s.n.* (NSW, PERTH); 8 June 2008, *M.N. Lyons 6068* (PERTH); 12 May 1972, *P.G. Wilson s.n.* (PERTH); 7 July 1973, *P.G. Wilson 11426* (PERTH).

*Phenology.* Flowers and fruits from February to July.

*Distribution and habitat.* Recorded in woodland, sometimes associated with pools on sandstone. Endemic to the Kimberley region and known only from the Bonaparte Archipelago on Augustus Island, Uwins Island, St Andrews Island (Lyons *et al.* 2013) and the adjacent mainland near Kunmunya Hill.

*Conservation status.* *Actinoschoenus quadricostatus* is listed by Jones (2014) as Priority One under Department of Parks and Wildlife Conservation Codes for Western Australian Flora, under the name *A. sp. C Kimberley Flora* (P.G. Wilson s.n. 12/5/72).




Figure 5. *Actinoschoenus quadricostatus*. A – plant base; B – culms; C – inflorescence; D – SEM of culm; E – SEM of glume; F – SEM of nut. Scale bars = 500  $\mu$ m (D); 1 mm (E, F). Images from *M.N. Lyons 6020*. Photographs by R.L. Barrett.

*Etymology.* The epithet is from the Latin *quadri-* (four-) and *costatus* (ribbed), in reference to the four-ribbed culms.

*Notes.* Morphologically similar to *A. arthrotyloides* but differing in having very slender, 4-ribbed culms (vs 5(6)-ribbed), 1–3 spikelets per head (vs (1)2–7), and with upper glumes glabrous or nearly so (vs minutely hairy distally on sides and midvein).

The vernacular name of Four-ribbed Actinoschoenus is suggested.

**Actinoschoenus ramosus** Rye, R.L.Barrett & M.D.Barrett, *sp. nov.*

*Type:* upper Lawley River, Lawley River National Park, Western Australia [precise locality withheld for conservation reasons], 12 March 2014, *R.L. Barrett* RLB 8988 (*holo:* PERTH 08614911; *iso:* BRI, DNA, K, NE, NSW).

*Actinoschoenus* sp. A, B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1037, Figure 310A (1992).

*Actinoschoenus* sp. A Kimberley Flora (C.R. Dunlop 5303), G. Paczkowska & A.R. Chapman, *West. Austral. Fl.: Descr. Cat.*, p. 41 (2000); Western Australian Herbarium, in *FloraBase*, <http://florabase.dpaw.wa.gov.au/> [accessed 1 March 2014].

*Illustration.* B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1038, Figure 310A (1992) [as *Actinoschoenus* sp. A].

*Perennial*, caespitose sedge, 0.3–0.6 m high. *Culms* 0.3–0.4 mm diam., lax, obtusely 3- or 4-ribbed, scabridulous immediately below inflorescence, elsewhere glabrous or nearly so. *Leaves* usually with short hairs on blade and orifice of sheath, glabrous or slightly shortly hairy below; uppermost leaf 45–85 mm long, with a blade 1–7 mm long. *Inflorescence* a loose simple or compound arrangement of 3 or more spikelets, the peduncle and each ray bearing 1 spikelet; primary rays 2–4, 3–8 mm long; secondary rays 0–2. *Bracts* 2 or 3, ciliate to almost glabrous on sheath, shortly hairy on blade, the longest bract basal, the basal bract 4–7 mm long, with a blade 3–5 mm long. *Spikelets* not very compressed, narrowly ovate in outline, 5.5–7 mm long, 1–1.5 mm wide. *Glumes* c. 6, 4-ranked to slightly spirally arranged, green on keel and ferruginous on sides, glabrous; upper glumes narrowly ovate, 4.5–5 mm long, usually slightly apiculate. *Stamens* 3; anther 3–4 mm long. *Style* undivided for 2–4 mm; base triangular in outline, c. 0.5 mm long, c. 0.3 mm wide; stigmatic branches 3, about as long as or longer than remainder of style. *Nut* with a stipe or basal contraction 0.3–0.5 mm long; body whitish, 1.3–1.4 mm long, 0.9–1.1 mm wide, transversely tuberculate-ribbed. C<sub>3</sub> photosynthetic pathway inferred from anatomy. (Figures 1H, 6)

*Diagnostic characters.* *Inflorescence* of 3 or more spikelets, with 2–4 primary rays and often also secondary rays, each bearing 1 spikelet, the basal bract longest. *Upper glumes* 4.5–5 mm long.

*Other specimens examined.* WESTERN AUSTRALIA: [localities withheld for conservation reasons] 25 Jan. 2007, *R.L. Barrett & M.D. Barrett* RLB 3860 (K, NSW, PERTH); 25 Jan. 2007, *R.L. Barrett & M.D. Barrett* RLB 3955 A (NSW, PERTH); 22 Apr. 2008, *R.L. Barrett & M.D. Barrett* RLB 4529 (PERTH); 15 Jan. 2010, *R.L. Barrett & M. Maier* RLB 6046 (NE, NSW, PERTH); 19 Jan. 2010, *R.L. Barrett, M. Maier & P. Kendrick* RLB 6263 (CANB, PERTH); 25 Jan. 2010, *R.L. Barrett & M.D. Barrett* RLB 6431 (PERTH); 24 Feb. 1980, *C.R. Dunlop* 5303 (BRI, *n.v.*, CANB, DNA, NSW, NT, PERTH); 19 Jan. 1982, *K.F. Kenneally* 7796 (NSW, PERTH).




Figure 6. *Actinoschoenus ramosus*. A – habitat on sand over broken sandstone; B – lax habit; C – culms; D – openly branching inflorescence; E – spikelets showing style branches; F – spikelets showing anthers; G – SEM of culm; H – SEM of glume; I – SEM of nut. Scale bars = 500  $\mu\text{m}$  (G); 1 mm (H, I). Images from *R.L. Barrett* RLB 8988 (A–F) and *R.L. Barrett* RLB 3860 (G–I). Photographs by R.L. Barrett.

**Phenology.** Flowers and fruits recorded from January to February.

**Distribution and habitat.** Apparently endemic to the Kimberley region where it is known between the Lawley River, Mitchell Plateau and the Prince Regent River. Recorded in sand amongst sandstone rocks or in skeletal sand on sandstone in open woodland, often between boulders that offer a degree of shading; one record is ‘adjacent to a watercourse’.

**Conservation status.** Recently listed as Priority Three under Department of Parks and Wildlife Conservation Codes for Western Australian Flora (Western Australian Herbarium 1998–) under the name *A. sp.* A Kimberley Flora (C.R. Dunlop 5303).

**Etymology.** The epithet is from the Latin *ramosus* (branched) in reference to the openly branched inflorescence.

*Notes.* The morphology of *A. ramosus* is very distinct and it is easily separated from all other named species in the genus by the openly branched inflorescence, the other species all having spikelets more or less sessile in a head.

Collections from the vicinity of the East Alligator River, Northern Territory (e.g. *C.R. Dunlop* 4403; *R. Fensham* 872; *J. Russell-Smith* 8423 & *J. Brock*) known by the phrase name *Fimbristylis* sp. Deaf Adder Gorge (*C.R. Dunlop* 4403) at DNA are evidently very closely related to *A. ramosus*. Indeed, the DNA duplicate of *C.R. Dunlop* 5303 (the voucher specimen for *A. sp.* A Kimberley Flora, upon which *A. ramosus* is based), is determined there as *F. sp.* Deaf Adder Gorge, highlighting their close similarity. While specimens from both states are very similar in general appearance, the Northern Territory collections have smaller spikelets and fruits, and thinner, 3(4)-ribbed culms relative to the thicker, (3)4-ribbed culms of the Western Australian collections seen. The culm differences are illustrated in Figure 1H–K. The differences in spikelet, fruit and culm size may represent clinal variation within a single taxon or *F. sp.* Deaf Adder Gorge may be a distinct taxon; further study of *F. sp.* Deaf Adder Gorge is underway by one of us (KLC) and it is retained as separate here pending the outcome of that study.

The vernacular name of Soft Actinoschoenus is suggested.

### **Fimbristylis** Vahl

#### **Fimbristylis helicophylla** Rye, R.L.Barrett & M.D.Barrett, *sp. nov.*

*Type:* east of Theda Station Homestead, Western Australia [precise locality withheld for conservation reasons], 11 March 2014, *R.L. Barrett* RLB 8945 (*holo:* PERTH 08614881; *iso:* BM, BRI, CANB, DNA, GENT, MEL, NE, NSW, NY, PERTH).

*Fimbristylis* sp. G, B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1093, Figure 317R (1992).

*Fimbristylis* sp. G Kimberley Flora (A.C. Beauglehole 51810), G. Paczkowska & A.R. Chapman, *West. Austral. Fl.: Descr. Cat.*, p. 49 (2000); Western Australian Herbarium, in *FloraBase*, <http://florabase.dpaw.wa.gov.au/> [accessed 1 March 2014].

*Illustration.* B.L. Rye in J.R. Wheeler (ed.), *Fl. Kimberley Reg.*, p. 1076, Figure 317R (1992) [as *Fimbristylis* sp. G].

*Perennial* tufted sedge, 0.35–0.60 m high, with a short rhizome. *Culms* 1.0–1.4 mm wide, striate, glabrous. *Lowest leaves* initially distichous, usually 5–17 cm long, the sheath green to pale brown to dark red-brown, almost glabrous with just a few ciliate hairs at the apex, the blade 2.0–3.6 mm wide, twisted along its length, glabrous, or with scattered ciliate hairs on margins, margins pale, thickened. *Culm-sheathing leaves* 1 or 2 per culm, reduced to sheath, almost glabrous with just a few ciliate hairs at the membranous apex, 8.5–12 cm long. *Inflorescence* a cyme or compound cyme, with a central sessile spikelet and 1–4 pedunculate lateral spikelets; primary rays 1–3, erect to spreading, 15–30 mm long. *Bracts* usually 2, reduced to sheath, rather glume-like, with broad, membranous margins reaching apex, ciliate, the basal bract 5.3–6.2 mm long. *Spikelets* pale brown, ovoid or narrowly obloid-ellipsoid to almost globular, 6.2–10.4(–12.1) mm long, 3.1–6.1 mm diam., many-flowered, obtuse; axis not prominently winged. *Glumes* spirally arranged, with a prominent green and red-brown keel and red-brown-spotted, broad, membranous margins, oblong-elliptic to almost circular, 4–5 mm long, ciliate,

emarginate. *Stamens* 3; filament white, 4–6 mm long, anther 2.0–2.3 mm long. *Style* triquetrous, glabrous except on base or slightly scabridulous on angles, undivided for *c.* 3 mm; base prominently enlarged, triangular to broadly triangular in outline, 0.3–0.4 mm wide, ciliate on each angle and with larger, patent to retrorse hairs at base, the largest *c.* 0.4 mm long; stigmatic branches 3, shorter than remainder of style. *Nut* very shortly stipitate, brown to black, trigonous with convex faces, subterete, broadly obovate in outline, 0.8–1.0 mm long, 0.7–0.8 mm wide, tuberculate. (Figure 7)

*Diagnostic characters.* *Perennial sedge.* *Culms* 1.0–1.4 mm wide, striate. *Leaves* with blade 2.0–3.6 mm wide, twisted along its length, margins pale, thickened. *Inflorescence* a cyme or compound cyme, with a central sessile spikelet and 1–4 pedunculate lateral spikelets. *Spikelets* ovoid to almost globular, 6.2–10.4(–12.1) mm long. *Glumes* spirally arranged, with red-brown-spotted, broad, membranous margins, 4–5 mm long, ciliate. *Nut* trigonous, 0.8–1.0 mm long, 0.7–0.8 mm wide, tuberculate.




Figure 7. *Fimbristylis helicophylla*. A – habitat, B – habit, C – leafy rosettes with twisted leaves, D – branched inflorescence, E – spikelets showing anthers, F – spikelet showing style branches, G – SEM of leaf, H – SEM of glume, I – SEM of nut. Scale bars = 500  $\mu$ m (G), 1 mm (H), 200  $\mu$ m (I). Images from R.L. Barrett RLB 8945. Photographs by R.L. Barrett



*Other specimens examined.* WESTERN AUSTRALIA: [localities withheld for conservation reasons] 22 Feb. 2005, *M.D. Barrett* MDB 1643 B (PERTH); 25 Mar. 2010, *M.D. Barrett & R.L. Barrett* MDB 2781 (PERTH); 4 Dec. 1994, *R.L. Barrett & M.D. Barrett* RLB 1009 (PERTH); 16 Feb. 2006, *R.L. Barrett & M.D. Barrett* RLB 3117 (NE, NSW, PERTH); 6 June 2012, *R.L. Barrett* RLB 7687 (DNA, NE, NSW, PERTH); 31 May 1976, *A.C. Beaglehole* 51810 (NT, PERTH).

*Phenology.* Flowers and fruits recorded for March to June.

*Distribution and habitat.* Possibly endemic to the Kimberley region, known from Theda and Doongan Stations on the Kalumburu–Gibb River road, and south to Prince Regent National Park and the Harding Range.

*Conservation status.* *Fimbristylis helicophylla* is listed by Jones (2014) as Priority One under Department of Parks and Wildlife Conservation Codes for Western Australian Flora under the name *F.* sp. G Kimberley Flora (A.C. Beaglehole 51810).

*Etymology.* The epithet is from the Greek *helicus* (spirally twisted) and *-phyllus* (leaved), in reference to the leaves of this species which are distinctly twisted along their length.

*Notes.* Superficially similar to *F. lanceolata* C.B. Clarke, with which it grows on Theda Station, in having broad leaves, but readily distinguished when fertile by the relatively short, ovoid to almost globular spikelets (6.2–10.4(–12.1) vs 10–35 mm long) with pale brown glumes (vs mid-brown). *Fimbristylis helicophylla* has broad spikelets on rather curved branches like *F. rara* R.Br. and similar species placed in sect. *Leptocladae* Ohwi by Kern (1974). *Fimbristylis rara* is an annual with mid-brown (vs pale brown) spikelets, which are often shorter (2.5–7(–13) vs 6.2–10.4(–12.1) mm long). The broad, twisted leaves of *F. helicophylla* are unusual in this section.

The vernacular name of Twisted Leaf *Fimbristylis* is recommended.

### Acknowledgements

Preparation of this paper was funded by the Botanic Gardens and Parks Authority as part of the 50<sup>th</sup> anniversary celebrations of the Western Australian Botanic Garden at Kings Park. Butch and Robyn Maher are thanked for their company and for sharing local knowledge on field trips over the past 15 years. Paul Doughty and the Western Australian Museum are thanked for allowing us to participate in biological surveys in the Prince Regent River Reserve in 2007 and 2010. Michi Maier (Biota), Peter Kendrick (DPaW) and Butch Maher (Fitzroy Helicopters) are thanked for assistance with collecting species in the north Kimberley. Peter and Pat Lacy are thanked for their generous hospitality at Mt Elizabeth Station. Traditional owners from the Kalumburu area are thanked for their assistance with fieldwork and for permission to access their traditional lands. Rick and Ann Jane and Bushtrack Safaris provided valuable logistical support for many research trips into the Prince Regent River area. Mike Lyons is thanked for finding his recent collections of *A. quadricostatus* to serve as the type. Terry Macfarlane, Juliet Wege, Kelly Shepherd, Ryonen Butcher, Kevin Thiele and curation staff at the Western Australian Herbarium are thanked for their support. Directors and staff at CANB, DNA, MEL, NE, NSW and PERTH are thanked for allowing access to their collections.

Some of the fieldwork associated with field studies of these species in 2010 was supported by a grant to the Western Australian Museum by Alcoa of Australia for the Alcoa Frog Watch programme and

a personal donation from Harry Butler. The Botanic Gardens and Parks Authority, Mark Webb and Kingsley Dixon are thanked for funding fieldwork in March 2010 and March 2014. Australian Biological Resources Study is thanked for funding to RLB for fieldwork in March 2014 on *Cleome* that provided additional opportunities to collect Cyperaceae, and to KLC for her PhD studies. Cecelia Myers and Dunkeld Pastoral supported fieldwork on Theda Station between 2005 and 2012. Additional support came from the 2010 JobsFund grant to Zoos South Australia.

## References

- Bruhl, J.J., Stone, N.E. & Hattersley, P.W. (1987). C<sub>4</sub> acid decarboxylation enzymes and anatomy in sedges (Cyperaceae): first record of NAD-malic enzyme species. *Australian Journal of Plant Physiology* 14: 719–728.
- Bruhl, J.J. & Wilson, K.L. (2008). Towards a comprehensive survey of C<sub>3</sub> and C<sub>4</sub> photosynthetic pathways in Cyperaceae. *Aliso* 23: 99–148.
- CHAH (Council of Heads of Australasian Herbaria (2007–). *Australian Plant Census (APC)*, IBIS database. Centre for Australian National Biodiversity Research, Canberra. <http://www.chah.gov.au/apc/index.html> [accessed 1 July 2015].
- Fitzgerald, W.V. (1918). The botany of the Kimberleys, north-west Australia. *Journal and Proceedings of the Royal Society of Western Australia* 3: 102–224.
- Hattersley, P.W. & Watson, L. (1975). Anatomical parameters for predicting photosynthetic pathways of grass leaves: the ‘maximum lateral cell count’ and the ‘maximum cells distant count’. *Phytomorphology* 25: 325–333.
- Hinchliff, C.E. & Roalson, E.H. (2013). Using supermatrices for phylogenetic inquiry: an example using the sedges. *Systematic Biology* 62: 205–219.
- Jones, A. (2014). *Threatened and Priority Flora list for Western Australia*. (Department of Parks and Wildlife: Kensington, Western Australia.)
- Kern, J.H. (1974). Cyperaceae. In: van Steenis, C.G.G.J. (ed.) *Flora Malesiana Series I*. Vol. 7(3). pp. 435–753. (Wolters-Noordhoff Publishing: Groningen.)
- Latz, P.K. (1990). Taxonomic studies of *Fimbristylis* (Cyperaceae) in northern Australia. *Nuytsia* 7: 161–182.
- Lyons, M.N., Keighery, G.J., Gibson, L.A. & Handasyde, T. (2013). Flora and vegetation communities of selected islands off the Kimberley coast of Western Australia. In: Gibson, L.A., Yates, S. & Doughty, P. (eds.) *Biodiversity values on selected Kimberley islands, Australia. Records of the Western Australian Museum Supplement No. 81*. pp. 205–243. (Western Australian Museum: Welshpool, Western Australia.)
- Rye, B.L. (1992). Cyperaceae. In: Wheeler, J.R. (ed.) *Flora of the Kimberley region*. pp. 1035–1107. (Conservation and Land Management: Perth, Western Australia.)
- Western Australian Herbarium (1998–). *FloraBase—the Western Australian Flora*. Department of Parks and Wildlife. <http://florabase.dpaw.wa.gov.au/> [accessed 1 March 2014].